



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

LLNL-TR-415408

Science Based Stockpile Stewardship, Uncertainty Quantification, and Surrogate Reactions

M. A. Stoyer, D. P. McNabb, J. T. Burke, L. A.
Bernstein

August 6, 2009

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

This work performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

Science Based Stockpile Stewardship, Uncertainty Quantification, and Surrogate Reactions

Mark A. Stoyer, Dennis P. McNabb, Jason T. Burke, Lee A. Bernstein

Stewardship of this nation's nuclear weapons is predicated on developing a fundamental scientific understanding of the physics and chemistry required to describe weapon performance without the need to resort to underground nuclear testing and to predict expected future performance as a result of intended or unintended modifications. In order to construct more reliable models, underground nuclear test data is being reanalyzed in novel ways. To improve the interpretation of these experiments with quantified uncertainties, improved nuclear data is required.

As an example, the thermonuclear yield of a device was often inferred through the use of radiochemical detectors. Conversion of the detector activations observed to thermonuclear yield was accomplished through explosion code calculations (models) and a good set of nuclear reaction cross-sections. Because of the unique high-fluence environment of an exploding nuclear weapon, many reactions occurred on radioactive nuclides, for which only theoretically calculated cross-sections are available. Surrogate nuclear reactions at STARS/LIBERACE offer the opportunity to measure cross-sections on unstable nuclei and thus improve the quality of the nuclear reaction cross-section sets.

One radiochemical detector that was loaded in devices was mono-isotopic yttrium (^{89}Y). Nuclear reactions produced ^{87}Y and ^{88}Y which could be quantified post-shot as a ratio of $^{87}\text{Y}/^{88}\text{Y}$. The yttrium cross-section set from 1988 is shown in Figure 1(a) and contains approximately 62 cross-sections interconnecting the yttrium nuclides. The 6 experimentally measured cross-sections are shown in Figure 1(b). Any measurement of cross-sections on ^{87}Y or ^{88}Y would improve the quality of the cross-section set. A recent re-evaluation of the yttrium cross-section set was performed with many more calculated reaction cross-sections included [1].




Fig. 1: The radiochemical cross-section set used for yttrium (a) and those cross-sections where experimental data exists (b). Reactions affecting the production or destruction of ^{87}Y and ^{88}Y are of interest to the Science Based Stockpile Stewardship program. Note several cross-sections are only measured at 14 MeV and only one cross-section on a radioactive target ($^{88}\text{Y}(n,2n)^{87}\text{Y}$) has been measured.

[1] R.D. Hoffman, et al., UCRL-TR-222275 (2006).